

Silvotator: Understanding diffeomorphisms by how they act on curves

① Introduction: the standard curve complex

Let S be a surface. The simplicial complex $\mathcal{C}(S)$ is given by:

- vertices = isotopy classes of essential simple closed curves in S (essential = not homotopic to a point, a puncture or a boundary component)

- edges: there is an edge b/w a and b if $i(a, b) = 0$ (intersection \emptyset)

$\mathcal{C}(S)$ is the flag complex of this graph (\rightarrow add all the simplices generated by these edges)

$\mathbb{R}k \text{ Mod}(S)$ acts on $\mathcal{C}(S)$ by simplicial transfo

Thm 0 If $3g + m \geq 5$ then $\mathcal{C}(S_{g,m})$ is connected

(Today: case $g \geq 1$)

ex $\mathcal{C}(S_{1,0})$: definitely not connected \Rightarrow discrete!

It is more interesting to make an edge $a - b$ if $i(a, b) = 1$ instead

Thm [Masur - Minsky 85] $\mathcal{C}(S)$ is Gromov δ -hyperbolic

(It was recently shown that δ does not depend on S)

($\delta =$ in a triangle \triangle \cup δ -nhl of 2 edges contains the third)

$\mathcal{N}(S)$ = subgraph generated by non-separating simple closed curves

② Modified version $\hat{\mathcal{N}}(S)$

$\hat{\mathcal{N}}(S)$: vertices are the same as $\mathcal{N}(S)$

there is an edge $a - b$ if $i(a, b) = 1$

lemma (Lickorish) $\mathcal{N}(S_{g,m})$ is connected for $g \geq 1$ and $m \geq 0$

proof Let a, b be vertices of $\hat{\mathcal{N}}(S)$. We need to find $c_0 = a, c_1, \dots, c_k = b$ such that $i(c_i, c_{i+1}) = 1$.



We construct this by induction on $i(a, b)$

* $i(a, b) = 1 \rightarrow$ nothing to do

* suppose $i(a, b) = n \geq 2$, assume the result proved for lower values of i
we look for γ s.t. $c(a, \gamma), c(b, \gamma) < n$

Suppose $i(a, b) = m > 0$, assume we know g for some values of g
 we look for γ s.t. $c(a, \gamma), c(b, \gamma) < m$

choose representatives α, β in minimal position, i.e. $\#(\alpha \cap \beta) = m$
 & choose two consecutive intersection points P, Q on β





Let $\gamma_1 = \alpha_1 \cup l, \gamma_2 = \alpha_2 \cup l$.

\hookrightarrow one of these must be a non-separating closed curve:

three regions ①②③. If both γ_1 and γ_2 were separating,
 then $① \cap ③ = \emptyset, ② \cap ③ = \emptyset$
 $\Rightarrow (① \cup ②) \cap ③ = \emptyset \Rightarrow \alpha$ is separating, contradiction


Pick γ_i the non-separating one & push it a bit inwards to get γ'_i




Moreover, since γ'_i is not separating, it is essential

* If $i(a, b) = 0$: two cases

• If $a \cup b$ is separating:



• If $a \cup b$ is not separating




pf of thm 0: Let $a, b \in \mathcal{C}(S)$

* If $a \cup b$ is not separating. From the lemma $\exists a = c, c \cup b$

17

* If a, b are not separating. From the lemma, $\exists a = c_0, \dots, c_k = b$ st $i(c_i, c_{i+1}) = 1$. Let V_i be a regular nhd of $c_i \cup c_{i+1}$, then ∂V_i is a sec st $i(\partial V_i, c_i) = 0, i(\partial V_i, c_{i+1}) = 0$



Moreover, if ∂V_i was not essential, then $S = S_{1,1}$
 \rightarrow forbidden

\rightarrow get a path $a = c_0, \partial V_0, c_1, \partial V_1, \dots, c_k = b$ in $\mathcal{C}(S)$

* If a is separating, $S \setminus a = S' \sqcup S''$. Both pieces have genus > 0 (b/c a is essential). Let a' be a non-separating curve in S'
 \rightarrow then $i(a, a') = 0$. We can thus assume that a is not separating

Thm If $G \subset X$, X : connected graph, the action is transitive on vertices and on pairs of vertices related by an edge

Let $v, w \in \text{Sk}_0(X)$ st $v - w \in \text{Sk}_1(X)$. Let $h \in G$ st $h \cdot w = v$
 Then G is generated by $h \cup \text{Stab}(v)$

cor $\text{Mod}(S_g)$ is finitely generated